

Table 1C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Demographic Indicators ⁽²⁾

Page: 1
Run Date: 7 / 12 / 11

Total Records	Number	%	Source of Data	Number	%
Complete Records	25,507	75.5	WIC	33,789	100.0
Prenatal Records	2,043	6.0	Maternal Health	0	0.0
Postpartum Records	6,239	18.5	Other	0	0.0
Total Records Accepted	33,789		Total	33,789	100.0
			<i>Excluded Unknown: 0</i>	<i>0.0</i>	

Race/Ethnic Distribution	Number	%	Age Distribution	Number	%	Education Distribution	Number	%
White, Not Hispanic	18,046	53.4	< 15 Years	82	0.2	< High School	9,167	29.2
Black, Not Hispanic	6,432	19.0	15 - 17 Years	1,711	5.1	High School	14,136	45.0
Hispanic	6,371	18.9	18 - 19 Years	4,210	12.5	> High School	8,076	25.7
American Indian/Alaska Native	611	1.8	20 - 29 Years	20,780	61.5	Total	31,379	100.0
Asian/Pacific Islander	1,584	4.7	30 - 39 Years	6,561	19.4			
Multiple Races	745	2.2	>= 40 Years	431	1.3			
All Other	0	0.0	Total	33,775	100.0			
Total	33,789	100.0	<i>Excluded Unknown: 14</i>	<i>0.0</i>		<i>Excluded Unknown: 2,410</i>	<i>7.1</i>	
<i>Excluded Unknown: 0</i>	<i>0.0</i>							

% Poverty Level	Number	%	Program Participation ⁽³⁾	Number	%	Migrant Status	Number	%
0 - 50	7,369	26.8	WIC	27,550	100.0	Yes	38	0.1
51 - 100	5,812	21.2	SNAP	27,205	49.1	No	33,751	99.9
101 - 130	2,789	10.2	Medicaid	27,477	34.6	Total	33,789	100.0
131 - 150	1,463	5.3	TANF	27,477	1.3			
151 - 185	1,816	6.6	<i>Excluded Unknown:</i>					
186 - 200	1	0.0	WIC:	0	0.0			
> 200	0	0.0	SNAP:	345	1.3			
Adjunctive Eligibility	8,227	29.9	Medicaid:	73	0.3			
Total	27,477	100.0	TANF:	73	0.3	<i>Excluded Unknown: 0</i>	<i>0.0</i>	
<i>Excluded Unknown: 73</i>	<i>0.3</i>							

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Prenatal program participation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Health Indicators

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Prepregnancy BMI (3)				
Underweight	33,415	3.5	1,226,875	4.5
Normal Weight	33,415	39.9	1,226,875	42.5
Overweight	33,415	26.0	1,226,875	25.9
Obese	33,415	30.6	1,226,875	27.0
Maternal Weight Gain (4)				
Less Than Ideal	26,656	22.8	917,032	21.2
Ideal	26,656	28.0	917,032	30.6
Greater Than Ideal	26,656	49.2	917,032	48.2
Anemia (Low Hb/Hct) (5)				
1st Trimester	11,951	7.1	436,474	7.7
2nd Trimester	9,887	11.9	375,216	12.2
3rd Trimester	5,309	32.1	189,944	33.9
Postpartum	27,098	23.9	412,633	28.8
Parity (6)				
0	31,792	39.2	1,108,534	41.3
1	31,792	28.1	1,108,534	26.7
2	31,792	17.5	1,108,534	16.3
≥ 3	31,792	15.2	1,108,534	15.6
Interpregnancy Interval				
< 6 Months	16,363	9.9	614,765	12.0
6 - <18 Months	16,363	28.6	614,765	26.7
≥18 Months	16,363	61.5	614,765	61.3
Gestational Diabetes	30,743	5.2	389,161	5.0
Hypertension During Pregnancy	30,743	5.8	394,368	6.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

(6) Number of times a woman has given birth to an infant 20 or more weeks gestation.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Health Indicators

Page: 3
Run Date: 7 / 12 / 11

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Medical Care				
1st Trimester	31,228	87.2	527,184	82.2
No Care	31,228	2.5	527,184	3.1
WIC Enrollment				
1st Trimester	33,671	35.8	1,213,982	34.1
2nd Trimester	33,671	29.8	1,213,982	34.7
3rd Trimester	33,671	15.9	1,213,982	17.5
Postpartum (3)	33,671	18.4	1,213,982	13.7
Prepregnancy Multivitamin Consumption (4)	32,690	18.5	360,369	18.7
Multivitamin Consumption During Pregnancy (5)	26,605	85.5	372,743	83.5
Smoking				
3 Months Prior to Pregnancy	32,048	34.5	1,040,568	22.9
During Pregnancy	27,483	21.9	895,533	13.8
Last 3 Months of Pregnancy	30,707	18.4	838,781	13.0
Postpartum	30,716	19.9	782,035	15.0
Smoking Changes (6)				
Quit Smoking by First Prenatal Visit	9,469	37.8	206,583	43.1
Quit Smoking by First Prenatal Visit and Stayed Off Cigarettes	8,468	32.8	156,709	36.7
Smoking in Household				
During Pregnancy	26,931	17.4	650,648	17.1
Postpartum	30,720	5.3	559,995	16.8
Drinking				
3 Months Prior to Pregnancy	31,926	12.6	1,040,654	8.3
Last 3 Months of Pregnancy	30,689	0.2	604,343	1.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who initially enrolled in WIC during the postpartum period.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

(6) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 2C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Health Indicators

Page: 4
Run Date: 7 / 12 / 11

Summary of Health Indicators (2)				
	Wisconsin Prevalence		National Prevalence (Prior Year)	
	Number	%	Number	%
Birthweight (3)				
Very Low	29,684	1.2	1,018,145	1.1
Low	29,684	6.8	1,018,145	7.1
Normal	29,684	85.0	1,018,145	85.1
High	29,684	7.1	1,018,145	6.7
Preterm (4)	30,784	9.9	1,017,333	11.1
Full term Low Birth Weight (5)	26,732	3.1	848,957	3.5
Multiple Births	31,565	1.3	1,095,174	1.4
Breastfeeding				
Ever Breastfed	30,131	70.4	997,136	66.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Very low birth weight (<1500 g), low birth weight (1500 - <2500 g), normal birth weight (2500 - <4000 g), high birth weight (>4000 g).

(4) <37 weeks gestation.

(5) 37 or more weeks gestation and <2500 g.



Table 3C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 5
Run Date: 7 / 12 / 11

Comparison of Racial and Ethnic Distribution by Contributor to Nation

Comparison of Racial and Ethnic Distribution ⁽²⁾								
	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
Wisconsin	33,789	53.4	19.0	18.9	1.8	4.7	2.2	0.0
Nation (Prior Year)	1,304,320	41.6	24.1	28.3	1.1	3.2	1.2	0.5

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 4C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Age Distribution by Contributor to Nation

Comparison of Age Distribution ⁽²⁾								
	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
Wisconsin	33,789	0.2	5.1	12.5	61.5	19.4	1.3	0.0
Nation (Prior Year)	1,304,320	0.3	5.8	12.3	59.4	20.4	1.5	0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Maternal Health Indicators by Contributor to Nation

Page: 7
Run Date: 7 / 12 / 11

Comparison of Maternal Health Indicators ⁽²⁾														
	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes During Pregnancy		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
Wisconsin	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9	30,743	5.2	30,743	5.8
Nation (Prior Year)	1,226,875	4.5	52.9	917,032	21.2	48.2	189,944	33.9	412,633	28.8	389,161	5.0	394,368	6.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 6C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 8
Run Date: 7 / 12 / 11

Comparison of Maternal Behavioral Indicators by Contributor to Nation

Comparison of Maternal Behavioral Indicators ⁽²⁾									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregpregnancy ⁽³⁾		During Pregnancy ⁽⁴⁾	
	Number	%	%	Number	%	Number	%	Number	%
Wisconsin	31,228	87.2	2.5	33,671	35.8	32,690	18.5	26,605	85.5
Nation (Prior Year)	527,184	82.2	3.1	1,213,982	34.1	360,369	18.7	372,743	83.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Comparison of Smoking Indicators by Contributor to Nation

Page: 9
Run Date: 7 / 12 / 11

Comparison of Smoking Indicators ⁽²⁾										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes ⁽³⁾	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum		
	Number	%	Number	%		Number	%	Number	%	
Wisconsin	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3
Nation (Prior Year)	1,040,568	22.9	838,781	13.0	156,709	36.7	650,648	17.1	559,995	16.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 8C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Comparison of Infant Health Indicators by Contributor to Nation

Page: 10
Run Date: 7 / 12 / 11

Comparison of Infant Health Indicators (2)									
	Birthweight			Preterm (5)	Full Term LBW (6)	Ever Breastfed			
	Low (3)		High (4)						
	Number	%	%						
Wisconsin	29,684	7.9	7.1	30,784	9.9	26,732	3.1	30,131	70.4
Nation (Prior Year)	1,018,145	8.2	6.7	1,017,333	11.1	848,957	3.5	997,136	66.7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 9C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Maternal Health Indicators by Race/Ethnicity, Age, or Education

Maternal Health Indicators by Race/Ethnicity, Age, or Education ⁽²⁾

	Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)				Gestational Diabetes		Hypertension During Pregnancy	
	Under- weight		Over- weight	< Ideal		> Ideal	3rd Trimester		Postpartum					
	Number	%	%	Number	%	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity														
White, Not Hispanic	17,871	3.9	54.3	14,056	19.3	53.1	2,867	23.5	14,248	16.7	16,345	4.9	16,345	6.0
Black, Not Hispanic	6,334	3.2	61.5	5,169	27.1	47.3	1,150	54.8	5,212	42.4	5,847	3.1	5,847	8.1
Hispanic	6,292	2.2	60.8	5,226	25.9	43.8	808	31.7	5,267	22.3	5,822	7.4	5,822	4.3
American Indian/Alaska Native	607	2.1	68.0	445	22.7	55.7	104	32.7	409	28.1	546	7.3	546	4.4
Asian/Pacific Islander	1,572	5.3	41.5	1,163	32.5	30.9	268	29.9	1,365	33.4	1,504	6.8	1,504	1.5
Multiple Races	739	3.7	58.3	597	24.0	50.3	112	28.6	597	25.1	679	5.0	679	5.7
All Other/Unknown	0	*	*	0	*	*	0	*	0	*	0	*	0	*
Total	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9	30,743	5.2	30,743	5.8
Age														
< 15 Years	82	*	*	66	*	*	16	*	63	*	74	*	74	*
15 - 17 Years	1,701	6.1	33.0	1,386	22.7	51.4	264	35.2	1,398	32.0	1,577	1.8	1,577	4.6
18 - 19 Years	4,158	6.0	42.2	3,321	20.4	51.9	648	39.5	3,307	29.7	3,814	1.3	3,814	5.6
20 - 29 Years	20,549	3.4	57.3	16,422	22.4	49.6	3,353	32.2	16,722	23.6	18,980	4.5	18,980	5.3
30 - 39 Years	6,489	1.4	69.3	5,152	25.4	45.7	965	26.3	5,299	19.6	5,941	10.0	5,941	7.7
>= 40 Years	422	0.7	71.8	300	33.0	41.0	61	*	309	16.5	344	18.9	344	11.3
Unknown	14	*	*	9	*	*	2	*	0	*	13	*	13	*
Total	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9	30,743	5.2	30,743	5.8
Education														
< High School	9,088	3.8	54.2	7,360	26.4	45.6	1,331	39.7	7,426	28.6	8,337	4.8	8,337	5.2
High School	14,012	3.8	56.9	11,342	21.9	50.4	2,309	32.0	11,330	24.0	12,948	4.9	12,948	5.4
> High School	8,008	2.3	59.1	6,397	19.7	51.8	1,350	25.0	6,502	18.5	7,354	5.9	7,354	7.4
Unknown	2,307	4.0	56.2	1,557	25.9	46.4	319	32.0	1,840	23.4	2,104	5.5	2,104	4.9
Total	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9	30,743	5.2	30,743	5.8

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 10C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education

Maternal Behavioral Indicators by Race/Ethnicity, Age, or Education (2)									
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester		No Care	1st Trimester		Pregnancy (3)		During Pregnancy (4)	
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	16,610	90.8	2.0	17,992	35.9	17,474	23.3	13,895	89.9
Black, Not Hispanic	5,923	84.0	1.6	6,405	30.4	6,238	9.5	4,964	83.2
Hispanic	5,951	82.8	4.5	6,341	43.5	6,157	16.5	5,336	80.5
American Indian/Alaska Native	555	84.1	2.5	607	38.7	602	15.0	526	85.6
Asian/Pacific Islander	1,503	77.5	3.3	1,581	25.0	1,491	9.8	1,286	67.6
Multiple Races	686	86.9	2.0	745	37.6	728	16.5	598	87.6
All Other/Unknown	0	*	*	0	*	0	*	0	*
Total	31,228	87.2	2.5	33,671	35.8	32,690	18.5	26,605	85.5
Age									
< 15 Years	75	*	*	81	*	79	*	71	*
15 - 17 Years	1,592	77.8	2.9	1,709	37.6	1,665	7.1	1,464	85.9
18 - 19 Years	3,865	86.0	2.7	4,196	40.4	4,084	8.9	3,595	84.1
20 - 29 Years	19,232	88.1	2.3	20,708	34.8	20,121	17.7	16,180	85.7
30 - 39 Years	6,065	88.2	2.6	6,532	35.9	6,328	29.6	4,972	86.4
>= 40 Years	385	81.6	3.4	431	32.5	400	30.3	310	81.0
Unknown	14	*	*	14	*	13	*	13	*
Total	31,228	87.2	2.5	33,671	35.8	32,690	18.5	26,605	85.5
Education									
< High School	8,496	81.4	3.7	9,133	39.3	8,892	11.2	7,575	80.4
High School	13,143	88.1	2.2	14,080	36.3	13,772	16.9	11,358	87.0
> High School	7,443	92.1	1.6	8,058	33.2	7,872	29.4	6,066	90.8
Unknown	2,146	87.0	2.7	2,400	29.0	2,154	18.8	1,606	79.6
Total	31,228	87.2	2.5	33,671	35.8	32,690	18.5	26,605	85.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for behavioral indicators, and data with errors for behavioral and demographic indicators.

(3) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(4) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 11C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Smoking Indicators by Race/Ethnicity, Age, or Education

Smoking Indicators by Race/Ethnicity, Age, or Education ⁽²⁾										
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
Race/Ethnicity										
White, Not Hispanic	17,309	45.5	16,321	25.7	5,976	29.6	14,096	20.0	16,330	5.3
Black, Not Hispanic	5,906	23.8	5,840	12.7	1,087	35.6	5,010	22.1	5,844	8.8
Hispanic	6,018	14.7	5,821	5.0	690	53.5	5,393	7.7	5,820	2.3
American Indian/Alaska Native	592	61.0	545	33.0	295	34.9	529	27.0	545	6.1
Asian/Pacific Islander	1,515	12.1	1,502	5.3	150	40.0	1,292	5.2	1,502	2.5
Multiple Races	708	48.2	678	25.2	270	32.2	611	23.9	679	7.2
All Other/Unknown	0	*	0	*	0	*	0	*	0	*
Total	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3
Age										
< 15 Years	79	*	74	*	10	*	71	*	74	*
15 - 17 Years	1,616	28.1	1,577	12.3	396	45.5	1,474	33.0	1,576	11.5
18 - 19 Years	3,986	39.7	3,809	16.8	1,293	39.9	3,635	27.4	3,809	7.4
20 - 29 Years	19,692	36.4	18,954	19.9	5,472	31.5	16,389	15.4	18,964	4.8
30 - 39 Years	6,259	27.8	5,936	16.9	1,250	27.4	5,035	12.3	5,940	4.1
>= 40 Years	403	20.3	344	12.5	45	*	314	13.1	344	3.5
Unknown	13	*	13	*	2	*	13	*	13	*
Total	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3
Education										
< High School	8,614	35.7	8,330	21.1	2,450	29.2	7,645	22.1	8,332	7.5
High School	13,506	39.6	12,934	21.1	4,122	32.8	11,500	18.6	12,941	5.6
> High School	7,755	26.5	7,352	11.6	1,513	39.9	6,166	10.7	7,353	2.9
Unknown	2,173	26.4	2,091	15.4	383	26.4	1,620	12.4	2,094	3.8
Total	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for smoking indicators, and data with errors for smoking and demographic indicators.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 12C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Infant Health Indicators by Race/Ethnicity, Age, or Education

Infant Health Indicators by Race/Ethnicity, Age, or Education ⁽²⁾									
	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
Race/Ethnicity									
White, Not Hispanic	15,770	7.2	8.3	16,363	9.6	14,251	2.6	16,006	72.0
Black, Not Hispanic	5,598	12.6	3.4	5,846	13.1	4,861	5.2	5,686	57.1
Hispanic	5,685	5.6	7.7	5,840	8.1	5,202	2.3	5,771	83.6
American Indian/Alaska Native	530	4.7	12.5	540	8.0	486	1.4	536	66.2
Asian/Pacific Islander	1,446	7.9	3.0	1,512	7.9	1,335	3.7	1,464	53.8
Multiple Races	655	8.2	7.9	683	9.2	597	3.0	668	70.7
All Other/Unknown	0	*	*	0	*	0	*	0	*
Total	29,684	7.9	7.1	30,784	9.9	26,732	3.1	30,131	70.4
Age									
< 15 Years	67	*	*	73	*	55	*	68	*
15 - 17 Years	1,509	6.6	4.8	1,581	8.8	1,381	3.8	1,530	64.2
18 - 19 Years	3,699	7.1	5.1	3,837	9.0	3,360	3.5	3,747	67.7
20 - 29 Years	18,336	6.5	7.0	19,014	9.6	16,563	3.0	18,618	70.4
30 - 39 Years	5,732	7.2	9.0	5,920	11.3	5,076	2.9	5,822	73.9
>= 40 Years	328	10.4	12.2	346	12.7	285	3.5	333	72.7
Unknown	13	*	*	13	*	12	*	13	*
Total	29,684	7.9	7.1	30,784	9.9	26,732	3.1	30,131	70.4
Education									
< High School	8,000	7.1	5.9	8,346	10.0	7,208	3.7	8,109	65.1
High School	12,495	7.0	7.0	12,911	9.8	11,244	3.2	12,651	67.7
> High School	7,137	6.1	8.5	7,355	9.8	6,437	2.3	7,264	80.3
Unknown	2,052	6.2	7.7	2,172	9.9	1,843	2.6	2,107	73.2
Total	29,684	7.9	7.1	30,784	9.9	26,732	3.1	30,131	70.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data for health indicators, and data with errors for health and demographic indicators.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 13C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin

Maternal Weight Gain and Birth Outcomes by Select Health Indicators

Maternal Weight Gain and Birth Outcomes by Select Health Indicators (2)										
	Weight Gain (3)			Birth Outcomes						
	< Ideal > Ideal			Birthweight			Preterm (6)		Full Term LBW (7)	
				Low (4)		High (5)				
	Number	%	%	Number	%	%	Number	%	Number	%
Prepregnancy BMI										
Underweight	933	30.5	30.4	1,008	15.8	1.8	1,047	14.7	859	8.0
Normal Weight	10,751	25.3	40.9	11,806	8.3	4.9	12,218	9.7	10,657	3.5
Overweight	6,903	15.9	60.2	7,686	7.3	7.5	7,916	9.5	6,945	2.6
Obese	8,069	24.7	52.9	8,960	7.0	10.3	9,263	9.8	8,074	2.3
Weight Gain										
< Ideal	-	-	-	5,536	13.2	3.3	5,746	14.3	4,749	5.4
Ideal	-	-	-	7,173	8.3	4.5	7,382	10.3	6,422	3.2
> Ideal	-	-	-	12,608	5.3	10.1	13,014	7.4	11,655	2.0
Smoking During Pregnancy										
Smokers	4,633	23.6	49.5	5,144	10.6	4.7	5,327	10.9	4,587	4.9
Non-Smokers	16,961	22.6	49.3	18,965	6.3	7.9	19,433	8.4	17,367	2.5

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 14C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Racial and Ethnic Distribution

Page: 16
Run Date: 7 / 12 / 11

Summary of Trends in Racial and Ethnic Distribution ⁽²⁾								
Year	Number Included in Analysis	White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %
2010	33,789	53.4	19.0	18.9	1.8	4.7	2.2	0.0
2009	35,896	53.7	19.1	18.9	1.9	4.5	1.9	0.0
2008	35,549	53.7	19.1	19.2	1.7	4.5	1.8	0.0
2007	34,934	53.6	19.3	19.1	2.0	4.5	1.7	0.0
2006	31,077	52.1	18.9	19.5	2.0	4.2	1.6	1.7
2005	29,354	54.7	19.2	17.2	2.2	4.3	0.4	2.0
2004	33,697	57.6	19.4	16.4	2.2	4.2	0.0	0.2
2003	0	*	*	*	*	*	*	*
2002	28,924	58.6	19.7	15.7	2.2	3.8	0.0	0.0
2001	31,080	58.3	20.0	15.5	2.2	4.0	0.0	0.0
2000	38,155	59.3	20.8	13.6	2.1	4.2	0.0	0.0
1999	30,872	61.1	20.9	11.9	2.3	3.9	0.0	0.0
1998	30,319	61.9	21.4	10.5	2.1	4.1	0.0	0.0
1997	30,380	62.3	21.4	9.5	2.4	4.3	0.0	0.0
1996	12,251	84.5	4.2	5.3	1.0	5.1	0.0	0.0
1995	29,036	62.4	21.8	9.0	2.1	4.7	0.0	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 15C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Age Distribution

Page: 17
Run Date: 7 / 12 / 11

Summary of Trends in Age Distribution ⁽²⁾								
Year	Number Included in Analysis	< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %
2010	33,789	0.2	5.1	12.5	61.5	19.4	1.3	0.0
2009	35,896	0.3	5.5	13.0	61.8	18.3	1.2	0.0
2008	35,549	0.3	6.2	13.4	61.6	17.2	1.1	0.0
2007	34,934	0.3	6.6	13.8	61.8	16.2	1.2	0.1
2006	31,077	0.4	6.7	14.1	62.1	15.5	1.1	0.1
2005	29,354	0.5	6.7	14.0	62.0	15.7	1.2	0.0
2004	33,697	0.5	6.7	14.7	61.0	16.0	1.2	0.0
2003	0	*	*	*	*	*	*	*
2002	28,924	0.5	7.4	15.9	59.6	15.5	1.1	0.0
2001	31,080	0.3	6.6	14.6	60.6	16.8	1.2	0.0
2000	38,155	0.5	8.0	16.4	58.7	15.3	1.1	0.0
1999	30,872	0.6	8.6	16.8	57.7	15.3	1.1	0.0
1998	30,319	0.7	8.8	16.6	57.2	15.6	1.0	0.0
1997	30,380	0.8	9.5	16.0	56.8	15.9	1.0	0.0
1996	12,251	0.4	7.2	15.7	58.7	17.0	1.0	0.0
1995	29,036	1.0	9.7	15.9	56.2	16.1	1.1	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 16C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Maternal Health Indicators

Page: 18
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Pregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
2010	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9
2009	35,482	3.7	55.1	26,554	21.6	50.9	5,697	31.9	28,469	25.0
2008	35,138	3.7	53.9	24,071	20.6	51.5	5,983	32.5	29,357	24.8
2007	34,503	3.9	53.1	24,133	21.0	50.7	5,810	33.1	29,138	25.8
2006	30,733	4.4	53.0	21,617	21.9	50.5	4,588	29.4	24,349	24.9
2005	25,573	4.5	50.4	21,542	24.5	45.5	4,675	30.4	19,010	25.3
2004	29,546	4.7	50.1	26,190	23.9	45.9	5,082	33.2	16,237	26.1
2003	0	*	*	0	*	*	0	*	0	*
2002	25,042	5.1	48.2	21,724	23.0	46.1	4,619	31.0	13,207	25.7
2001	27,218	5.4	47.2	23,533	23.7	45.1	2,855	17.0	14,327	26.4
2000	32,882	5.8	45.8	28,239	24.6	44.8	5,363	26.1	14,148	26.4
1999	27,032	6.0	44.9	23,018	25.2	44.0	4,840	24.5	10,218	25.6
1998	26,849	5.7	44.6	23,617	24.4	43.7	4,515	29.7	9,529	23.2
1997	26,105	5.8	43.7	23,105	25.0	43.1	4,565	27.7	9,599	21.4
1996	10,183	6.3	41.5	9,004	25.7	41.7	1,775	22.3	3,636	18.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 16C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Maternal Health Indicators

Page: 19
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Health Indicators ⁽²⁾										
Year	Pregpregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
	Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
	Number	%	%	Number	%	%	Number	%	Number	%
1995	24,836	6.4	40.5	21,703	26.0	42.0	4,377	31.9	8,906	21.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 17C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Maternal Behavioral Indicators

Page: 20
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Behavioral Indicators ⁽²⁾					
Year	Medical Care			WIC Enrollment	
	1st Trimester		No Care	1st Trimester	
	Number	%	%	Number	%
2010	31,228	87.2	2.5	33,671	35.8
2009	33,210	85.6	2.8	35,778	34.1
2008	32,437	84.0	2.3	34,433	33.0
2007	31,285	83.3	2.3	33,775	33.7
2006	11,903	83.9	0.1	27,099	29.8
2005	22,622	74.9	12.5	28,229	33.2
2004	30,590	74.3	13.9	32,024	34.7
2003	0	*	*	0	*
2002	25,779	74.1	12.5	27,533	33.2
2001	27,652	74.3	10.7	30,274	0.0
2000	33,694	72.5	12.4	36,735	21.1
1999	26,697	71.1	13.0	29,540	23.2
1998	26,953	71.1	12.4	28,767	36.1
1997	27,168	71.3	12.2	28,118	34.4
1996	10,761	72.9	12.9	11,274	36.0
1995	25,376	70.1	13.3	25,878	32.0

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 18C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Smoking Indicators

Page: 21
Run Date: 7 / 12 / 11

Summary of Trends in Smoking Indicators (2)										
Year	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
	Number	%	Number	%	Number	%	Number	%	Number	%
2010	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3
2009	34,096	36.1	32,593	19.7	9,317	31.5	28,409	20.4	32,610	6.9
2008	33,969	36.6	32,190	20.4	9,493	30.9	28,619	21.6	32,204	7.9
2007	33,094	35.6	31,575	21.6	8,882	28.0	27,956	22.7	31,534	8.8
2006	20,837	33.1	28,206	21.2	6,146	26.5	14,138	25.1	20,724	11.0
2005	18,399	39.8	22,131	26.4	6,285	30.0	18,584	39.8	26,480	33.6
2004	26,351	40.8	29,526	23.7	9,094	29.3	26,414	41.6	30,700	39.6
2003	0	*	0	*	0	*	0	*	0	*
2002	22,548	42.2	24,910	24.5	6,872	32.9	22,642	43.4	25,745	41.5
2001	23,803	42.5	26,637	25.5	8,350	26.8	23,769	44.9	27,572	42.9
2000	30,624	42.8	31,613	25.6	10,461	27.6	30,764	46.6	33,784	44.9
1999	24,100	44.4	25,437	26.2	8,424	28.5	24,204	47.1	27,080	45.2
1998	29,369	43.2	6,139	25.4	212	26.9	24,115	47.0	6,326	42.3
1997	29,464	42.2	6,059	25.4	0	*	23,678	47.4	6,332	40.4
1996	11,796	45.3	2,505	27.3	0	*	9,450	51.1	2,622	43.1
1995	27,869	41.5	6,453	26.2	0	*	21,792	49.0	6,828	43.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 19C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Infant Health Indicators

Summary of Trends in Infant Health Indicators ⁽²⁾									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
2010	29,684	7.9	7.1	30,784	9.9	26,732	3.1	30,131	70.4
2009	31,539	7.8	7.8	32,660	9.7	28,463	3.1	32,009	69.0
2008	31,033	7.7	7.8	31,344	10.2	27,048	3.1	31,499	67.0
2007	30,255	7.8	7.8	30,697	10.4	26,286	3.2	30,747	65.8
2006	26,732	7.7	7.7	23,522	10.8	20,880	3.0	27,572	64.1
2005	24,726	7.8	8.2	24,324	11.1	21,339	2.9	24,996	62.3
2004	28,209	7.6	8.6	27,611	11.4	24,125	3.0	28,189	62.2
2003	0	*	*	0	*	0	*	0	*
2002	23,758	7.4	9.2	23,341	10.7	20,350	3.1	23,977	59.2
2001	25,501	7.4	9.2	12,367	27.9	8,615	8.5	25,708	57.2
2000	29,964	7.3	9.4	23,960	12.8	20,250	3.9	30,279	55.3
1999	24,383	7.5	9.4	20,706	15.5	17,201	3.6	23,396	51.1
1998	25,133	7.6	9.3	24,536	11.6	21,191	2.8	20,339	42.3
1997	24,451	7.6	9.5	24,093	11.2	20,374	3.1	21,352	43.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 19C
2010 Pregnancy Nutrition Surveillance (1)
Wisconsin
Summary of Trends in Infant Health Indicators

Page: 23
Run Date: 7 / 12 / 11

Summary of Trends in Infant Health Indicators ⁽²⁾									
Year	Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
	Low (3)		High (4)						
	Number	%	%	Number	%	Number	%	Number	%
1996	9,702	6.4	11.0	9,469	10.3	8,208	2.5	9,966	56.0
1995	23,260	7.3	9.6	21,362	11.2	18,699	3.1	23,216	50.1

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 24
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
White, Not Hispanic	2010	17,871	3.9	54.3	14,056	19.3	53.1	2,867	23.5	14,248	16.7
	2009	19,076	4.3	53.0	14,596	18.4	54.9	3,136	25.0	15,122	17.7
	2008	18,891	4.1	51.9	13,316	17.9	55.1	3,223	24.0	15,677	17.5
	2007	18,521	4.6	50.9	13,326	17.9	54.4	3,130	25.4	15,672	18.7
	2006	16,072	5.1	50.2	11,870	18.6	54.4	2,511	22.4	13,019	17.5
	2005	14,082	5.4	47.7	12,245	21.4	48.7	2,574	22.8	10,446	18.5
	2004	17,132	5.4	47.3	15,317	21.3	48.8	2,882	25.7	8,996	18.9
	2003	0	*	*	0	*	*	0	*	0	*
	2002	14,802	5.8	45.6	13,020	21.3	48.6	2,593	22.6	7,489	18.1
	2001	15,938	6.3	44.3	14,007	21.5	47.4	1,547	10.9	7,780	19.0
Black, Not Hispanic	2010	6,334	3.2	61.5	5,169	27.1	47.3	1,150	54.8	5,212	42.4
	2009	6,764	3.0	59.5	4,535	25.1	50.0	1,173	53.3	5,488	43.9
	2008	6,706	3.6	58.7	3,818	23.1	52.1	1,282	53.8	5,556	44.8
	2007	6,649	3.2	59.3	3,986	23.8	51.0	1,252	54.8	5,530	45.1
	2006	5,804	3.8	60.5	3,864	25.7	50.2	902	52.1	4,483	45.2
	2005	5,101	3.4	58.2	4,408	26.9	45.9	965	52.3	3,563	43.4
	2004	5,957	4.0	57.3	5,329	26.9	45.2	1,092	55.3	3,010	46.4
	2003	0	*	*	0	*	*	0	*	0	*
	2002	5,084	4.3	55.7	4,320	24.3	46.3	949	54.9	2,321	44.0
	2001	5,692	4.4	54.0	4,824	26.3	44.3	750	33.2	3,026	42.7

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 25
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Hispanic	2010	6,292	2.2	60.8	5,226	25.9	43.8	808	31.7	5,267	22.3
	2009	6,688	2.5	58.9	5,336	25.4	44.4	888	28.8	5,472	24.2
	2008	6,696	2.1	57.7	5,076	23.7	45.5	958	31.8	5,725	23.0
	2007	6,528	2.8	56.2	4,961	23.6	44.5	901	29.7	5,616	24.6
	2006	5,978	2.7	56.9	4,202	24.1	44.5	794	26.4	4,916	24.0
	2005	4,217	2.6	53.4	3,407	30.5	37.4	728	26.0	3,610	24.3
	2004	4,532	2.8	53.9	3,903	26.8	40.7	730	31.1	3,117	25.5
	2003	0	*	*	0	*	*	0	*	0	*
	2002	3,658	3.0	50.0	3,088	25.5	40.3	741	29.8	2,650	28.2
	2001	3,896	3.5	51.0	3,242	25.4	40.7	407	13.0	2,711	26.3
American Indian/Alaska Native	2010	607	2.1	68.0	445	22.7	55.7	104	32.7	409	28.1
	2009	673	1.2	64.8	506	21.3	49.8	91	*	499	26.3
	2008	619	3.9	63.5	450	19.3	51.1	89	*	477	26.6
	2007	683	1.8	60.6	473	21.1	55.0	106	37.7	526	32.5
	2006	619	3.1	62.7	420	22.9	51.2	67	*	466	25.8
	2005	579	2.1	62.0	511	22.1	52.3	85	*	431	28.3
	2004	670	2.8	61.0	582	22.2	51.4	95	*	379	24.8
	2003	0	*	*	0	*	*	0	*	0	*
	2002	575	4.0	59.7	499	20.6	52.7	81	*	210	35.2
	2001	634	2.4	57.6	553	21.7	51.0	73	*	220	33.6

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 26
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
Asian/Pacific Islander	2010	1,572	5.3	41.5	1,163	32.5	30.9	268	29.9	1,365	33.4
	2009	1,614	4.8	40.8	1,081	32.8	32.8	308	29.5	1,356	32.7
	2008	1,572	5.8	35.8	970	33.2	28.9	308	34.7	1,387	33.4
	2007	1,543	5.4	36.0	1,012	36.2	28.6	324	31.2	1,329	31.2
	2006	1,309	6.0	34.2	886	36.6	28.3	243	31.3	1,080	32.9
	2005	1,073	6.2	32.5	896	34.9	26.5	260	33.8	858	35.5
	2004	1,203	5.7	33.2	1,058	36.3	24.7	278	31.3	730	33.2
	2003	0	*	*	0	*	*	0	*	0	*
	2002	923	7.5	33.9	797	37.4	23.3	255	29.8	537	37.2
	2001	1,058	6.6	34.3	907	37.9	24.6	78	*	590	36.8
Multiple Races	2010	739	3.7	58.3	597	24.0	50.3	112	28.6	597	25.1
	2009	667	3.3	55.6	500	18.0	53.6	101	31.7	532	24.8
	2008	653	2.9	56.5	441	17.7	57.4	123	35.8	535	23.9
	2007	577	3.3	55.3	375	21.9	53.9	97	*	465	25.4
	2006	486	3.3	50.8	347	21.9	51.9	67	*	385	26.2
	2005	86	*	*	72	*	*	19	*	76	*
	2004	1	*	*	1	*	*	0	*	1	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 20C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 27
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Health Indicators by Race/Ethnicity

Summary of Trends in Maternal Health Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Prepregnancy BMI (3)			Weight Gain (4)			Anemia (Low Hb/Hct) (5)			
		Under-weight		Over-weight	< Ideal		> Ideal	3rd Trimester		Postpartum	
		Number	%	%	Number	%	%	Number	%	Number	%
All Other/Unknown	2010	0	*	*	0	*	*	0	*	0	*
	2009	0	*	*	0	*	*	0	*	0	*
	2008	1	*	*	0	*	*	0	*	0	*
	2007	2	*	*	0	*	*	0	*	0	*
	2006	465	5.6	48.6	28	*	*	4	*	0	*
	2005	435	6.7	45.7	3	*	*	44	*	26	*
	2004	51	*	*	0	*	*	5	*	4	*
	2003	0	*	*	0	*	*	0	*	0	*
	2002	0	*	*	0	*	*	0	*	0	*
	2001	0	*	*	0	*	*	0	*	0	*
Total All Race/Ethnic Groups	2010	33,415	3.5	56.6	26,656	22.8	49.2	5,309	32.1	27,098	23.9
	2009	35,482	3.7	55.1	26,554	21.6	50.9	5,697	31.9	28,469	25.0
	2008	35,138	3.7	53.9	24,071	20.6	51.5	5,983	32.5	29,357	24.8
	2007	34,503	3.9	53.1	24,133	21.0	50.7	5,810	33.1	29,138	25.8
	2006	30,733	4.4	53.0	21,617	21.9	50.5	4,588	29.4	24,349	24.9
	2005	25,573	4.5	50.4	21,542	24.5	45.5	4,675	30.4	19,010	25.3
	2004	29,546	4.7	50.1	26,190	23.9	45.9	5,082	33.2	16,237	26.1
	2003	0	*	*	0	*	*	0	*	0	*
	2002	25,042	5.1	48.2	21,724	23.0	46.1	4,619	31.0	13,207	25.7
	2001	27,218	5.4	47.2	23,533	23.7	45.1	2,855	17.0	14,327	26.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(5) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 28
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
White, Not Hispanic	2010	16,610	90.8	2.0	17,992	35.9
	2009	17,837	89.7	2.0	19,219	33.5
	2008	17,417	88.9	1.9	18,530	33.0
	2007	16,803	88.0	1.9	18,195	33.6
	2006	6,548	88.8	0.0	14,333	28.6
	2005	12,797	80.5	10.5	15,420	33.9
	2004	17,664	79.9	11.4	18,387	35.6
	2003	0	*	*	0	*
	2002	15,218	80.2	10.0	16,080	34.7
	2001	16,234	80.0	9.1	17,667	0.0
Black, Not Hispanic	2010	5,923	84.0	1.6	6,405	30.4
	2009	6,324	81.3	2.5	6,834	30.7
	2008	6,203	78.9	2.3	6,490	29.3
	2007	6,015	78.7	2.5	6,409	30.2
	2006	2,401	78.6	0.0	5,031	28.3
	2005	4,516	71.7	10.1	5,478	30.6
	2004	5,927	71.7	11.8	6,346	32.3
	2003	0	*	*	0	*
	2002	4,911	70.2	11.4	5,505	29.4
	2001	5,386	70.9	8.5	6,004	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 29
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Hispanic	2010	5,951	82.8	4.5	6,341	43.5
	2009	6,266	81.0	5.1	6,756	41.5
	2008	6,202	79.4	3.3	6,641	38.4
	2007	5,934	79.0	3.0	6,444	39.3
	2006	1,904	81.0	0.2	5,269	33.3
	2005	3,751	63.4	22.7	4,852	35.1
	2004	5,040	61.7	24.8	5,239	37.3
	2003	0	*	*	0	*
	2002	4,096	61.2	22.7	4,303	34.9
	2001	4,291	62.8	18.8	4,704	0.0
American Indian/Alaska Native	2010	555	84.1	2.5	607	38.7
	2009	621	82.8	3.2	674	40.1
	2008	550	81.6	3.5	586	36.7
	2007	598	75.8	2.5	656	39.2
	2006	294	70.7	0.0	519	36.8
	2005	520	78.7	4.8	616	42.4
	2004	660	72.7	10.8	695	44.0
	2003	0	*	*	0	*
	2002	550	75.3	8.9	593	40.6
	2001	596	73.8	8.2	672	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 30
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
Asian/Pacific Islander	2010	1,503	77.5	3.3	1,581	25.0
	2009	1,530	74.9	3.5	1,623	21.7
	2008	1,472	69.2	2.7	1,542	22.5
	2007	1,411	68.4	3.2	1,502	22.4
	2006	562	66.5	0.0	1,069	18.1
	2005	996	59.0	13.1	1,203	20.2
	2004	1,298	58.3	17.1	1,338	18.7
	2003	0	*	*	0	*
	2002	1,004	53.1	15.0	1,052	18.3
	2001	1,145	52.8	12.9	1,227	0.0
Multiple Races	2010	686	86.9	2.0	745	37.6
	2009	632	87.5	1.9	672	36.0
	2008	593	81.8	1.7	644	37.7
	2007	524	81.5	3.4	569	38.0
	2006	194	82.0	0.0	432	27.8
	2005	42	*	*	102	24.5
	2004	1	*	*	0	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 21C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 31
Run Date: 7 / 12 / 11

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity

Summary of Trends in Maternal Behavioral Indicators by Race/Ethnicity ⁽²⁾						
Race/Ethnicity and Year		Medical Care			WIC Enrollment	
		1st Trimester		No Care	1st Trimester	
		Number	%	%	Number	%
All Other/Unknown	2010	0	*	*	0	*
	2009	0	*	*	0	*
	2008	0	*	*	0	*
	2007	0	*	*	0	*
	2006	0	*	*	446	65.0
	2005	0	*	*	558	44.6
	2004	0	*	*	19	*
	2003	0	*	*	0	*
	2002	0	*	*	0	*
	2001	0	*	*	0	*
Total All Race/Ethnic Groups	2010	31,228	87.2	2.5	33,671	35.8
	2009	33,210	85.6	2.8	35,778	34.1
	2008	32,437	84.0	2.3	34,433	33.0
	2007	31,285	83.3	2.3	33,775	33.7
	2006	11,903	83.9	0.1	27,099	29.8
	2005	22,622	74.9	12.5	28,229	33.2
	2004	30,590	74.3	13.9	32,024	34.7
	2003	0	*	*	0	*
	2002	25,779	74.1	12.5	27,533	33.2
	2001	27,652	74.3	10.7	30,274	0.0

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 22C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Smoking Indicators by Race/Ethnicity

Page: 32
Run Date: 7 / 12 / 11

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
White, Not Hispanic	2010	17,309	45.5	16,321	25.7	5,976	29.6	14,096	20.0	16,330	5.3
	2009	18,512	48.4	17,536	27.8	6,773	28.7	14,880	23.7	17,551	7.2
	2008	18,401	49.3	17,301	29.0	6,896	28.2	14,914	25.2	17,312	8.0
	2007	17,872	48.3	16,939	30.7	6,495	25.7	14,581	26.5	16,933	9.2
	2006	10,856	46.6	15,000	30.3	4,548	24.4	7,370	29.6	11,021	11.6
	2005	10,309	54.4	12,936	35.2	4,844	28.5	10,372	49.5	14,813	41.5
	2004	14,915	54.7	17,033	31.6	6,932	28.0	14,929	51.1	17,671	48.3
	2003	0	*	0	*	0	*	0	*	0	*
	2002	12,939	56.8	14,651	32.3	5,350	32.2	12,961	52.8	15,175	49.9
	2001	13,605	56.4	15,583	33.5	6,407	26.1	13,591	54.5	16,177	51.3
Black, Not Hispanic	2010	5,906	23.8	5,840	12.7	1,087	35.6	5,010	22.1	5,844	8.8
	2009	6,370	24.5	6,197	12.8	1,117	35.3	5,350	26.0	6,197	10.3
	2008	6,417	25.4	6,123	14.0	1,212	33.3	5,518	27.1	6,123	12.5
	2007	6,318	24.2	6,023	13.9	1,116	30.4	5,408	27.1	6,015	13.3
	2006	3,921	22.6	5,379	14.2	745	27.2	2,598	29.4	3,935	16.3
	2005	3,604	25.8	4,190	18.7	762	27.4	3,668	34.3	5,199	30.0
	2004	5,228	27.6	5,669	17.8	1,187	27.0	5,266	36.8	5,954	35.1
	2003	0	*	0	*	0	*	0	*	0	*
	2002	4,477	28.7	4,741	19.1	866	27.4	4,511	38.4	4,922	36.6
	2001	4,818	30.2	5,151	19.8	1,126	24.1	4,821	40.0	5,366	38.4

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Smoking Indicators by Race/Ethnicity

Page: 33
Run Date: 7 / 12 / 11

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Hispanic	2010	6,018	14.7	5,821	5.0	690	53.5	5,393	7.7	5,820	2.3
	2009	6,362	14.4	6,101	5.5	731	48.4	5,697	8.1	6,102	2.6
	2008	6,386	14.4	6,153	5.1	734	50.3	5,732	9.0	6,152	3.4
	2007	6,227	13.4	6,026	6.1	653	45.3	5,576	10.2	6,011	3.8
	2006	4,269	12.0	5,552	5.3	455	44.0	2,912	12.7	4,051	5.2
	2005	3,228	11.7	3,526	6.1	319	52.7	3,274	17.2	4,610	14.4
	2004	4,483	13.5	4,896	5.7	516	51.2	4,492	18.1	5,077	17.3
	2003	0	*	0	*	0	*	0	*	0	*
	2002	3,722	12.1	3,986	4.8	343	56.0	3,742	20.1	4,081	19.3
	2001	3,771	13.4	4,174	5.9	410	45.9	3,756	21.5	4,276	20.9
American Indian/Alaska Native	2010	592	61.0	545	33.0	295	34.9	529	27.0	545	6.1
	2009	647	58.4	611	32.9	301	28.9	567	33.9	612	13.1
	2008	601	59.9	545	32.1	278	29.9	529	30.8	547	9.7
	2007	648	55.1	612	32.5	284	26.1	580	32.9	605	11.7
	2006	437	50.8	562	33.8	196	24.5	302	37.4	425	11.3
	2005	453	68.2	509	43.4	271	32.5	455	58.9	578	50.7
	2004	628	65.1	628	41.6	345	26.1	629	61.7	663	58.4
	2003	0	*	0	*	0	*	0	*	0	*
	2002	532	64.3	530	40.6	247	30.8	546	59.2	553	57.9
	2001	582	70.3	584	45.5	341	22.9	588	63.4	604	62.1

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Smoking Indicators by Race/Ethnicity

Page: 34
Run Date: 7 / 12 / 11

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2010	1,515	12.1	1,502	5.3	150	40.0	1,292	5.2	1,502	2.5
	2009	1,564	10.7	1,532	4.0	138	48.6	1,356	4.5	1,532	2.3
	2008	1,528	9.5	1,482	4.6	112	35.7	1,353	6.6	1,483	3.3
	2007	1,467	9.9	1,451	5.2	115	47.8	1,312	7.5	1,448	3.9
	2006	1,002	8.4	1,251	3.8	75	*	714	8.0	945	5.7
	2005	799	12.6	930	6.5	87	*	803	21.0	1,180	17.5
	2004	1,097	12.0	1,299	5.1	114	45.6	1,098	21.1	1,334	20.0
	2003	0	*	0	*	0	*	0	*	0	*
	2002	878	9.7	1,002	4.9	66	*	882	21.3	1,014	20.4
	2001	1,027	8.1	1,145	3.6	66	*	1,013	16.5	1,149	16.5
Multiple Races	2010	708	48.2	678	25.2	270	32.2	611	23.9	679	7.2
	2009	641	49.0	616	25.8	257	35.0	559	28.4	616	9.9
	2008	636	49.8	586	25.3	261	37.5	573	29.1	587	9.9
	2007	561	49.0	524	30.3	219	25.6	498	32.1	522	9.4
	2006	346	41.9	458	25.3	127	26.0	236	27.1	343	16.3
	2005	6	*	40	*	2	*	12	*	100	22.0
	2004	0	*	1	*	0	*	0	*	1	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
	2001	0	*	0	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 22C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin
Summary of Trends in Smoking Indicators by Race/Ethnicity

Page: 35
Run Date: 7 / 12 / 11

Summary of Trends in Smoking Indicators by Race/Ethnicity ⁽²⁾											
Race/Ethnicity and Year		Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (3)		Smoking in Household			
		3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
		Number	%	Number	%	Number	%	Number	%	Number	%
All Other/Unknown	2010	0	*	0	*	0	*	0	*	0	*
	2009	0	*	0	*	0	*	0	*	0	*
	2008	0	*	0	*	0	*	0	*	0	*
	2007	1	*	0	*	0	*	1	*	0	*
	2006	6	*	4	*	0	*	6	*	4	*
	2005	0	*	0	*	0	*	0	*	0	*
	2004	0	*	0	*	0	*	0	*	0	*
	2003	0	*	0	*	0	*	0	*	0	*
	2002	0	*	0	*	0	*	0	*	0	*
2001	0	*	0	*	0	*	0	*	0	*	
Total All Race/Ethnic Groups	2010	32,048	34.5	30,707	18.4	8,468	32.8	26,931	17.4	30,720	5.3
	2009	34,096	36.1	32,593	19.7	9,317	31.5	28,409	20.4	32,610	6.9
	2008	33,969	36.6	32,190	20.4	9,493	30.9	28,619	21.6	32,204	7.9
	2007	33,094	35.6	31,575	21.6	8,882	28.0	27,956	22.7	31,534	8.8
	2006	20,837	33.1	28,206	21.2	6,146	26.5	14,138	25.1	20,724	11.0
	2005	18,399	39.8	22,131	26.4	6,285	30.0	18,584	39.8	26,480	33.6
	2004	26,351	40.8	29,526	23.7	9,094	29.3	26,414	41.6	30,700	39.6
	2003	0	*	0	*	0	*	0	*	0	*
	2002	22,548	42.2	24,910	24.5	6,872	32.9	22,642	43.4	25,745	41.5
	2001	23,803	42.5	26,637	25.5	8,350	26.8	23,769	44.9	27,572	42.9

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 23C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 36
Run Date: 7 / 12 / 11

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
White, Not Hispanic	2010	15,770	7.2	8.3	16,363	9.6	14,251	2.6	16,006	72.0
	2009	16,963	6.9	8.7	17,558	9.1	15,412	2.6	17,202	70.1
	2008	16,660	7.3	9.1	16,874	9.7	14,628	2.9	16,932	67.4
	2007	16,266	7.1	8.9	16,549	10.0	14,278	2.8	16,505	65.9
	2006	14,311	7.3	8.7	12,778	10.0	11,460	3.0	14,678	64.1
	2005	13,898	7.1	9.1	13,584	10.4	12,048	2.5	13,937	62.5
	2004	16,293	7.1	9.5	15,806	10.9	13,884	2.8	16,277	62.9
	2003	0	*	*	0	*	0	*	0	*
	2002	14,008	6.7	10.5	13,660	10.1	12,037	2.8	14,083	60.2
	2001	14,948	6.9	10.4	7,333	32.1	4,824	8.0	15,017	59.3
Black, Not Hispanic	2010	5,598	12.6	3.4	5,846	13.1	4,861	5.2	5,686	57.1
	2009	5,959	12.6	4.1	6,208	13.0	5,175	5.4	6,026	55.6
	2008	5,826	11.5	3.8	5,863	12.4	4,873	5.0	5,890	54.0
	2007	5,650	12.2	4.2	5,752	13.0	4,674	5.5	5,740	52.3
	2006	4,937	11.5	4.2	4,262	14.0	3,638	4.7	5,219	49.7
	2005	4,735	11.9	5.0	4,712	14.4	3,982	4.6	4,839	48.6
	2004	5,441	11.3	5.2	5,469	14.6	4,602	4.9	5,390	46.4
	2003	0	*	*	0	*	0	*	0	*
	2002	4,464	11.4	4.8	4,519	13.3	3,760	4.8	4,579	44.1
	2001	4,878	11.1	4.9	2,408	19.7	1,843	11.6	4,971	39.8

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 37
Run Date: 7 / 12 / 11

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Hispanic	2010	5,685	5.6	7.7	5,840	8.1	5,202	2.3	5,771	83.6
	2009	5,941	6.0	8.9	6,127	8.3	5,434	2.3	6,077	83.2
	2008	6,015	5.5	8.7	6,066	8.7	5,373	2.0	6,123	83.2
	2007	5,871	5.8	8.8	5,902	9.4	5,170	2.2	6,000	83.0
	2006	5,322	5.7	8.9	4,661	10.3	4,150	1.8	5,453	81.8
	2005	4,299	5.7	9.0	4,217	9.9	3,762	2.5	4,349	81.4
	2004	4,660	5.1	9.3	4,558	9.1	4,098	2.0	4,688	81.4
	2003	0	*	*	0	*	0	*	0	*
	2002	3,817	5.6	9.5	3,726	9.4	3,312	2.7	3,862	77.6
	2001	4,024	5.3	9.9	1,794	26.9	1,272	6.8	4,068	75.5
American Indian/Alaska Native	2010	530	4.7	12.5	540	8.0	486	1.4	536	66.2
	2009	591	3.4	14.2	612	8.5	541	1.1	596	66.9
	2008	518	5.6	12.7	512	13.1	422	0.9	522	62.1
	2007	572	5.8	14.3	580	11.6	486	3.1	582	59.1
	2006	534	6.2	12.9	452	9.5	406	1.7	549	65.8
	2005	540	6.3	16.5	531	11.5	464	1.9	532	62.0
	2004	588	5.4	14.3	563	13.5	477	1.5	577	60.8
	2003	0	*	*	0	*	0	*	0	*
	2002	509	4.1	15.5	480	11.7	418	1.2	481	61.5
	2001	539	4.3	16.7	221	19.5	170	7.1	525	59.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 38
Run Date: 7 / 12 / 11

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
Asian/Pacific Islander	2010	1,446	7.9	3.0	1,512	7.9	1,335	3.7	1,464	53.8
	2009	1,494	6.4	4.4	1,540	7.5	1,378	3.0	1,506	52.3
	2008	1,445	7.0	3.5	1,456	10.4	1,261	2.8	1,458	47.0
	2007	1,398	6.8	3.5	1,406	8.3	1,237	3.5	1,416	49.6
	2006	1,189	5.9	4.7	982	8.1	896	2.7	1,218	44.8
	2005	1,141	7.2	4.3	1,104	9.6	991	3.2	1,148	45.6
	2004	1,226	7.0	5.6	1,215	9.9	1,064	3.0	1,254	49.5
	2003	0	*	*	0	*	0	*	0	*
	2002	960	7.3	5.1	956	11.9	823	3.9	972	42.2
	2001	1,112	5.6	5.2	611	16.4	506	7.3	1,127	38.6
Multiple Races	2010	655	8.2	7.9	683	9.2	597	3.0	668	70.7
	2009	591	10.7	7.6	615	11.5	523	5.0	602	70.3
	2008	569	8.3	8.1	573	12.0	491	2.6	574	67.4
	2007	498	7.8	7.0	508	9.4	441	3.4	504	65.3
	2006	439	10.0	5.9	387	14.5	330	4.2	452	64.6
	2005	95	*	*	93	*	81	*	100	68.0
	2004	0	*	*	0	*	0	*	0	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 23C
2010 Pregnancy Nutrition Surveillance ⁽¹⁾
Wisconsin

Page: 39
Run Date: 7 / 12 / 11

Summary of Trends in Infant Health Indicators by Race/Ethnicity

Summary of Trends in Infant Health Indicators by Race/Ethnicity ⁽²⁾										
Race/Ethnicity and Year		Birthweight			Preterm (5)		Full Term LBW (6)		Ever Breastfed	
		Low (3)		High (4)						
		Number	%	%	Number	%	Number	%	Number	%
All Other/Unknown	2010	0	*	*	0	*	0	*	0	*
	2009	0	*	*	0	*	0	*	0	*
	2008	0	*	*	0	*	0	*	0	*
	2007	0	*	*	0	*	0	*	0	*
	2006	0	*	*	0	*	0	*	3	*
	2005	18	*	*	83	*	11	*	91	*
	2004	1	*	*	0	*	0	*	3	*
	2003	0	*	*	0	*	0	*	0	*
	2002	0	*	*	0	*	0	*	0	*
	2001	0	*	*	0	*	0	*	0	*
Total All Race/Ethnic Groups	2010	29,684	7.9	7.1	30,784	9.9	26,732	3.1	30,131	70.4
	2009	31,539	7.8	7.8	32,660	9.7	28,463	3.1	32,009	69.0
	2008	31,033	7.7	7.8	31,344	10.2	27,048	3.1	31,499	67.0
	2007	30,255	7.8	7.8	30,697	10.4	26,286	3.2	30,747	65.8
	2006	26,732	7.7	7.7	23,522	10.8	20,880	3.0	27,572	64.1
	2005	24,726	7.8	8.2	24,324	11.1	21,339	2.9	24,996	62.3
	2004	28,209	7.6	8.6	27,611	11.4	24,125	3.0	28,189	62.2
	2003	0	*	*	0	*	0	*	0	*
	2002	23,758	7.4	9.2	23,341	10.7	20,350	3.1	23,977	59.2
	2001	25,501	7.4	9.2	12,367	27.9	8,615	8.5	25,708	57.2

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

(3) <2500 g.

(4) >4000 g.

(5) <37 weeks gestation.

(6) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

